

UC Irvine

UC Irvine Previously Published Works

Title

Draft genome sequences of five new strains of methylophilaceae isolated from lake washington sediment.

Permalink

<https://escholarship.org/uc/item/3b66z0pw>

Journal

Genome announcements, 3(1)

ISSN

2169-8287

Authors

McTaggart, Tami L
Benuska, Gabrielle
Shapiro, Nicole
et al.

Publication Date

2015-02-01

DOI

10.1128/genomea.01511-14

Peer reviewed

Draft Genome Sequences of Five New Strains of *Methylophilaceae* Isolated from Lake Washington Sediment

Tami L. McTaggart,^{a*} Gabrielle Benuska,^a Nicole Shapiro,^b Tanja Woyke,^b Ludmila Chistoserdova^a

Department of Chemical Engineering, University of Washington, Seattle, Washington, USA^a; DOE Joint Genome Institute, Walnut Creek, California, USA^b

* Present address: Tami L. McTaggart, University of Irvine, Irvine, California, USA.

We sequenced the genomes of five new *Methylophilaceae* strains isolated from Lake Washington sediment. We used the new sequences to sort these new strains into specific *Methylophilaceae* ecotypes, including one novel ecotype. The new genomes expand the known diversity of *Methylophilaceae* and provide new models for studying the ecology of methylotrophy.

Received 16 December 2014 Accepted 22 December 2014 Published 5 February 2015

Citation McTaggart TL, Benuska G, Shapiro N, Woyke T, Chistoserdova L. 2015. Draft genome sequences of five new strains of *Methylophilaceae* isolated from Lake Washington sediment. *Genome Announc* 3(1):e01511-14. doi:10.1128/genomeA.01511-14.

Copyright © 2015 McTaggart et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ludmila Chistoserdova, milachis@u.washington.edu.

We recently described the diversity of *Methylophilaceae* from Lake Washington through an analysis of the genomes and the phenotypes of 11 strains, separating them into eight distinct ecotypes: *Methylotenera mobilis* JLW8, *M. mobilis* 13, *Methylotenera versatilis* 301, *Methylovorus glucosotrophus*, *Methylophilaceae* 7, *Methylophilaceae* 11, *Methylophilus methylotrophus* Brown, and *M. methylotrophus* White (1). Here, we report the draft genome sequences of five additional strains from Lake Washington sediment, isolated using the techniques described by Beck et al. (1). Strain 7 was isolated from the 2011 sample (1), strain G11 was isolated from the 2013 sample (2), strain L2L1 was isolated from long-term methane enrichment (2), and strains Q8 and N17 were isolated from short-term methane enrichment (2).

The draft genomes were generated at the Department of Energy (DOE) Joint Genome Institute (JGI) using Pacific Biosciences (PacBio) sequencing technology (3) and assembled using HGAP (version: 2.2.0.p1) (4), as part of the JGI sequencing pipeline (<http://www.jgi.doe.gov>). The genome statistics are shown in Table 1.

Strain Q8 is most similar to *M. methylotrophus* sp. strain 1, and strain L2L1 is most similar to *M. mobilis* 13 (Table 1). The main difference in metabolic potential is the lack of the gene for nitrous

oxide reductase in the genome of strain L2L1. Strain N17 is most similar to *Methylophilaceae* sp. strain 11 (Table 1). Strain 7 does not have close relatives; at the 16S rRNA gene identity level, it is related most closely to strain G11 and less related to the *Methylophilaceae* 7 ecotype. However, in terms of genome-genome similarity, this strain is more related to *Methylophilaceae* 7 (Table 1). Thus, we ascribed this strain to the *Methylophilaceae* 7 ecotype. Phenotypically, strain 7 is similar to other strains within this ecotype. However, it has a somewhat enhanced methylotrophy potential, encoding an additional XoxF enzyme and a true Fae that is missing from strains 73s and 79 (1). The protein encoded by this gene is most related to a Fae in the *M. glucosotrophus* SIP3-4 genome (83%) (5), suggesting a horizontal transfer event. At 2,503 Mb, this genome is the smallest so far for Lake Washington *Methylophilaceae*.

Strain G11 is the most novel strain of the set. In terms of 16S rRNA gene identity, it is most similar to *M. versatilis* 301, while in terms of genome-genome similarity, it is somewhat more related to *M. mobilis* JLW8 (Table 1). Thus, we identify this strain as a novel ecotype within Lake Washington *Methylophilaceae*. Like strains JLW8 and 301, G11 does not encode calcium-containing (MxaFI) methanol dehydrogenase but encodes three XoxF-type

TABLE 1 Strains described, closest relatives, genome statistics, accession numbers, and variable metabolic modules

Strain	Closest relative	% 16S rRNA gene identity	% proteins with >80% identity	Total no. of base pairs	NCBI accession no.	Presence of ^a :					
						MxaFI	MADH	NMGP	NarABC	NirK	NorBD
Q8	<i>M. methylotrophus</i> 1	100	86.4	2,900,053	JUHE00000000	+	+	—	—	—	—
L2L1	<i>M. mobilis</i> 13	99.6	79.3	2,641,989	JQMG00000000	+	—	—	+	+	+
N17	<i>Methylophilaceae</i> 11	99.9	88.5	2,753,666	JUGE01000001	+	—	—	—	—	—
7	<i>Methylophilaceae</i> 7	97.3	69.1	2,503,174	JUGF00000000	+	+	—	—	—	—
	<i>Methylophilaceae</i> G11	97.6	21.3	2,503,174	JUGF00000000	+	+	—	—	—	—
G11	<i>M. versatilis</i> 301	98.5	21.2	2,545,099	JUHH01000001	—	+	+	—	+	+
	<i>M. mobilis</i> JLW8	97.5	37.0	2,545,099	JUHH01000001	—	+	+	—	+	+

^aMxaFI, calcium-containing methanol dehydrogenase; MADH, methylamine dehydrogenase; NMGP, *N*-methylglutamate pathway; NarABC, NirK, dissimilatory nitrate and nitrite reductases; NorBD, nitric oxide reductase.

proteins. Like strain JLW8, it encodes methylamine dehydrogenase, and like strain 301, it encodes the *N*-methylglutamate pathway for methylamine oxidation (1).

The availability of these new genomes further expands the diversity of *Methylophilaceae* from a single ecological niche and further points toward unexpected metabolic flexibility within this taxon. The new strains described here also expand the repertoire of model organisms that can be used to study the ecology of methylotrophy.

Nucleotide sequence accession numbers. The sequences have been deposited in GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation (grant MCB-0950183) and by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under award no. DE-SC-0010556. The work conducted by the U.S. Department of Energy Joint Genome Institute was supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231.

REFERENCES

1. Beck DAC, McTaggart TL, Setboonsarng U, Vorobev A, Kalyuzhnaya MG, Ivanova N, Goodwin L, Woyke T, Lidstrom ME, Chistoserdova L. 2014. The expanded diversity of *Methylophilaceae* from Lake Washington through cultivation and genomic sequencing of novel ecotypes. *PLoS One* 9:e102458. <http://dx.doi.org/10.1371/journal.pone.0102458>.
2. Oshkin I, Beck DAC, Lamb AE, Tchesnokova V, Benuska G, McTaggart TL, Kalyuzhnaya MG, Dedysh S, Lidstrom ME, Chistoserdova L. 2014. Methane fed microcosms show differential community dynamics and pinpoint specific taxa involved in communal response. *ISME J*, in press. <http://dx.doi.org/10.1038/ismej.2014.203>.
3. Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, Zhao P, Zhong F, Korlach J, Turner S. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <http://dx.doi.org/10.1126/science.1162986>.
4. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
5. Lapidus A, Clum A, LaButti K, Kalyuzhnaya MG, Lim S, Beck DAC, Glavina del Rio T, Nolan M, Mavromatis K, Huntemann M, Lucas S, Lidstrom ME, Ivanova N, Chistoserdova L. 2011. Genomes of three methylotrophs from a single niche reveal the genetic and metabolic divergence of the *Methylophilaceae*. *J Bacteriol* 193:3757–3764. <http://dx.doi.org/10.1128/JB.00404-11>.